

Package ‘flowFitExampleData’

March 29, 2021

Type Package

Title Example data for the flowFit package

Version 1.26.0

Date 2013-03-04

Author davide Rambaldi

Maintainer Davide Rambaldi <davide.rambaldi@gmail.com>

Description Two dataset that can be used to run examples from the flowFit vignette and examples

License Artistic-2.0

biocViews FlowCytometryData

Depends R (>= 2.12.2), flowCore

Imports methods

PackageStatus Deprecated

git_url <https://git.bioconductor.org/packages/flowFitExampleData>

git_branch RELEASE_3_12

git_last_commit dd51ba5

git_last_commit_date 2020-10-27

Date/Publication 2021-03-29

R topics documented:

PKH26data	1
QuahAndParish	2
Index	4

PKH26data *genFitting: example data with PKH26 stain*

Description

example data with PKH26 dye. Two samples: NONPROL (parent population) and PROL (proliferating population).

Usage

```
data(PKH26data)
```

Format

The format is an object of class `flowSet` with 2 `flowFrame`

QuahAndParish

Example dataset from: New and improved methods for measuring lymphocyte proliferation in vitro and in vivo using CFSE-like fluorescent dyes (Benjamin J C Quah and Christopher R Parish, 2012)

Description

Detection of lymphocyte division by carboxyfluorescein diacetate succinimidyl ester (CFSE), Cell Trace Violet (CTV) and Cell Proliferation Dye eFluor 670 (CPD) in vitro.

Usage

```
data(QuahAndParish)
```

Format

The format is an object of class `flowSet` with 4 objects of class `flowFrame`

1. Fig 2a All CD4 T Nonstim.fcs Control sample including non-activated cells (non-dividing) labelled with CFSE, CPD and CTV
2. Fig 2a CFSE CD4 T Stim.fcs CD4 T cells stained with CFSE
3. Fig 2a CPD CD4 T Stim.fcs CD4 T cells stained with CPD
4. Fig 2a CTV CD4 T Stim.fcs CD4 T cells stained with CTV

The phenodata lists:

Filename The filename

SampleType The sample type (Nonstim or Stim)

Stain Stain type

CellType Cell type

Details

This QuahAndParish dataset represents the measurements of CD4 T cells division by CFSE, CTV and CPD in vitro. Spleen cells from B6 mice were labelled with $10\mu M$ CFSE, CTV and/or CPD and cultured for 4 days in the presence of a range of polyclonal stimuli that activate T and B cells. Viable CD4+ cells were discriminated using specific antibody staining. The dataset represent the measurements used in figure 2a (CD4+ population) from the paper: New and improved methods for measuring lymphocyte proliferation in vitro and in vivo using CFSE-like fluorescent dyes (Benjamin J C Quah and Christopher R Parish, 2012).

References

1. Benjamin J.C. Quah, Christopher R. Parish, New and improved methods for measuring lymphocyte proliferation in vitro and in vivo using CFSE-like fluorescent dyes, *Journal of Immunological Methods*, Volume 379, Issues 1-2, 31 May 2012, Pages 1-14, ISSN 0022-1759, 10.1016/j.jim.2012.02.012.

Index

* datasets

PKH26data, [1](#)

QuahAndParish, [2](#)

flowFrame, [2](#)

flowSet, [2](#)

PKH26data, [1](#)

QuahAndParish, [2](#)