

# Package ‘tissueTreg’

September 23, 2025

**Title** TWGBS and RNA-seq data from tissue T regulatory cells from mice

**Version** 1.29.0

**Author** Charles Imbusch [aut, cre],  
Michael Delacher [aut],  
Markus Feuerer [aut],  
Benedikt Brors [aut]

**Maintainer** Charles Imbusch <c.imbusch@dkfz.de>

**Description** The package provides ready to use epigenomes (obtained from TWGBS) and transcriptomes (RNA-seq) from various tissues as obtained in the study (Delacher and Imbusch 2017, PMID: 28783152).  
Regulatory T cells (Treg cells) perform two distinct functions: they maintain self-tolerance, and they support organ homeostasis by differentiating into specialized tissue Treg cells. The underlying dataset characterises the epigenetic and transcriptomic modifications for specialized tissue Treg cells.

**Depends** R (>= 3.5)

**License** GPL (>= 2)

**Encoding** UTF-8

**LazyData** true

**Suggests** BiocStyle, knitr, rmarkdown, testthat, ExperimentHub, bsseq, SummarizedExperiment, ggplot2, reshape2

**VignetteBuilder** knitr

**biocViews** ExperimentData, Tissue, Mus\_musculus\_Data, SequencingData, RNASeqData

**URL** <https://github.com/cimbusch/tissueTreg>

**RoxygenNote** 6.0.1

**git\_url** <https://git.bioconductor.org/packages/tissueTreg>

**git\_branch** devel

**git\_last\_commit** b245bce

**git\_last\_commit\_date** 2025-04-15

**Repository** Bioconductor 3.22

**Date/Publication** 2025-09-23

## Contents

|                      |          |
|----------------------|----------|
| tissueTreg . . . . . | 2        |
| <b>Index</b>         | <b>3</b> |

---

|            |                                                                            |
|------------|----------------------------------------------------------------------------|
| tissueTreg | <i>Epigenomes and transcriptomes of tissue resident regulatory T cells</i> |
|------------|----------------------------------------------------------------------------|

---

### Description

The package provides ready to use epigenomes (obtained from TWGBS) and transcriptomes (RNA-seq) from various tissues. Regulatory T cells (Treg cells) perform two distinct functions: they maintain self-tolerance, and they support organ homeostasis by differentiating into specialized tissue Treg cells. The underlying dataset characterises the epigenetic and transcriptomic modifications for specialized tissue Treg cells.

### Source

Delacher, M, Imbusch, CD, Weichenhan, D, Breiling, A, Hotz-Wagenblatt, A, Träger, U, Hofer, AC, Kägebein, D, Wang, Q, Frauhammer, F, Mallm, JP, Bauer, K, Herrmann, C, Lang, PA, Brors, B, Plass, C, Feuerer, M (2017). Genome-wide DNA-methylation landscape defines specialization of regulatory T cells in tissues. Nat. Immunol., 18, 10:1160-1172.

### Examples

```
eh <- ExperimentHub::ExperimentHub()

# RNA-seq RPKM data:
se_rpkms <- eh[["EH1074"]]

# Whole genome bisulfite sequencing data as bsseq objects:
tregs_per_sample <- eh[["EH1072"]]
tregs_per_tissue <- eh[["EH1073"]]
```

# Index

\* **datasets**

tissueTreg, [2](#)

tissueTreg, [2](#)